

Measles

Chief Health Officer Alert

Status:	Active
Date issued:	22 May 2017
Issued by:	Dr Brett Sutton, Deputy Chief Health Officer, Victoria
Issued to:	General Practitioners and Hospital Emergency Departments

Key messages

- **There is a confirmed case of** measles in a person who was infectious whilst on flight GA716 from Jakarta to Melbourne, departing Jakarta at 23.55 on Friday May 12 and arriving at Tullamarine at 09.20 on Saturday May 13.
- **The case has travelled internationally recently** and acquired the infection in Indonesia.
- **Be alert** for measles in patients presenting with a fever at rash onset, particularly associated with international travel.
- **Isolate suspected cases** to minimise the risk of transmission within your department/practice.
- **Notify** the Communicable Disease Prevention and Control Section at the Department of Health and Human Services on 1300 651 160 of suspected cases immediately.
- **Take blood for measles serology** in all suspected cases.
- **Discuss whether to take nose and throat swabs** for PCR with the Department if your suspicion for measles is high. Approval is required prior to PCR testing at the reference laboratory. PCR testing for measles does not attract a Medicare rebate.

What is the issue?

There is a confirmed case of measles in a person who was infectious whilst on an international flight and in the Tullamarine airport.

- Friday 12 May - Saturday May 13 Flight GA 716 (Garuda Indonesia) from Jakarta to Melbourne
- Saturday 13 May - Tullamarine airport from 9.20am - 12pm.

The illness was acquired in Indonesia.

Anyone who presents with signs and symptoms compatible with measles should be tested and notified to the Department. There should be an especially high index of suspicion if they have attended any of the areas stated above and are unvaccinated or partially vaccinated for measles.

Who is at risk?

Children or adults born during or since 1966 who do not have documented evidence of receiving two doses of a measles-containing vaccine or do not have documented evidence of immunity are considered to be susceptible to measles. People who are immunocompromised are also at risk.

Symptoms and transmission

Clinical features of measles include prodromal fever, a cough, conjunctivitis and coryza. Individuals, especially children, are typically unwell.

The most important clinical predictors are the following features:

- generalised, maculopapular rash, usually lasting three or more days, AND
- fever (at least 38°C, if measured) present at the time of rash onset, AND
- cough, coryza or conjunctivitis.

Measles is transmitted by airborne droplets and direct contact with discharges from respiratory mucous membranes of infected persons and less commonly, by articles freshly soiled with nose and throat secretions.

Measles is highly infectious and can persist in the environment for up to two hours.

The incubation period is variable and averages 10 days (range: 7 -18 days) from exposure to the onset of fever, with an average of 14 days from exposure to the onset of rash. The infectious period of patients with measles is roughly five days before, to four days after, the appearance of the rash.

Use the most appropriate tests for diagnosis depending on timing of symptoms and presentation:

- Take blood for serological confirmation in all suspected cases. If a patient has measles, IgM is reliably positive if the rash has been present for three or more days. IgG in the absence of IgM indicates the patient is protected and means measles is unlikely.
- Nose and throat swabs for PCR diagnosis are best for early diagnosis (including prior to rash); you must contact the Department prior to taking swabs to gain approval for these to be tested at the Victorian Infectious Diseases Reference Laboratory. PCR testing for measles does not attract a Medicare rebate.

This picture is typical of rash on the face. This is a rash on day three in a young boy.



Picture courtesy of U.S. Centers for Disease Control and Prevention

Recommendations

- Be alert for new measles cases – ensure all staff, especially triage nurses, have a high index of suspicion for patients presenting with a febrile rash or unwell following recent international travel.
- Notify suspected cases immediately to the Communicable Disease Prevention and Control Section via telephone on 1300 651 160 (24 hours).
- Take blood for serological confirmation.
- Call the department to discuss the need for PCR diagnosis.

- To minimise the risk of measles transmission within your department/practice:
 - avoid keeping patients with a febrile rash illness in shared waiting areas
 - give the suspected case a single use mask and isolate them, until a measles diagnosis can be excluded
 - leave vacant all consultation rooms used in the assessment of patients with suspected measles for at least 30 minutes after the consultation.
- Seek advice from the Department of Health and Human Services Communicable Disease Prevention and Control Section regarding:
 - the management of susceptible hospital or clinic contacts
 - prevention of measles in susceptible contacts.
- On advice, follow up all persons who attended the emergency department or clinic at the same time as a case and for 30 minutes after the visit. These people are considered to be exposed to the measles virus.
- Check your staff vaccination records.
- Earlier outbreaks have affected health care workers, including some who have not been involved in the direct care of measles cases and have only been in the same ward, clinic, or department as a case. All staff born during or since 1966 should have documentation of two doses of measles-containing vaccine, or laboratory-confirmed evidence of past measles infection.

More information

Clinical information

- [The Australian Immunisation Handbook: 10th edition, 2013](http://www.immunise.health.gov.au/internet/immunise/publishing.nsf/Content/Handbook10-home~handbook10part4~handbook10-4-9)
<http://www.immunise.health.gov.au/internet/immunise/publishing.nsf/Content/Handbook10-home~handbook10part4~handbook10-4-9>
- [The Blue Book – Guidelines for the control of infectious diseases](https://www2.health.vic.gov.au/public-health/infectious-diseases/disease-information-advice/measles)
<https://www2.health.vic.gov.au/public-health/infectious-diseases/disease-information-advice/measles>

Consumer information

- [Better Health Channel – Measles](http://www.betterhealth.vic.gov.au/bhcv2/bhcarticles.nsf/pages/Measles?open)
<http://www.betterhealth.vic.gov.au/bhcv2/bhcarticles.nsf/pages/Measles?open>

Contacts

For further information please contact the Communicable Disease Prevention and Control section at the Department of Health and Human Services on 1300 651 160 (24 hours).



Dr Brett Sutton
Deputy Chief Health Officer

Subscribe now!

Keep informed of emergencies affecting the health sector and critical public health issues impacting your work. [Subscribe now](https://www2.health.vic.gov.au/newsletters) <https://www2.health.vic.gov.au/newsletters> to information including Chief Health Officer Alerts and emergency advice from the Department of Health and Human Services.

Share this



Authorised and published by the Victorian Government, 1 Treasury Place, Melbourne.

© State of Victoria, Department of Health and Human Services, March 2017.

Available at [Health alerts and advisories](https://www2.health.vic.gov.au/about/news-and-events/healthalerts) <<https://www2.health.vic.gov.au/about/news-and-events/healthalerts>>