

Opioid Risk Tool

Date: _____

Patient Name: _____

| Mark each box that applies: | Female (Score) | Male (Score) |
|---|--|--|
| Family history of substance abuse <ul style="list-style-type: none"> • Alcohol • Illegal drugs • Prescription drugs | <input type="checkbox"/> 1 <input type="checkbox"/> 2 <input type="checkbox"/> 4 | <input type="checkbox"/> 3 <input type="checkbox"/> 3 <input type="checkbox"/> 4 |
| Personal history of substance abuse <ul style="list-style-type: none"> • Alcohol • Illegal drugs • Prescription drugs | <input type="checkbox"/> 3 <input type="checkbox"/> 4 <input type="checkbox"/> 5 | <input type="checkbox"/> 3 <input type="checkbox"/> 4 <input type="checkbox"/> 5 |
| Age (mark box if between 16-45 years) | <input type="checkbox"/> 1 | <input type="checkbox"/> 1 |
| History of preadolescent sexual abuse | <input type="checkbox"/> 3 | <input type="checkbox"/> 0 |
| Psychological disease <ul style="list-style-type: none"> • Attention deficit disorder, obsessive compulsive disorder, bipolar, schizophrenia • Depression | <input type="checkbox"/> 2 <input type="checkbox"/> 1 | <input type="checkbox"/> 2 <input type="checkbox"/> 1 |
| Total | _____ | _____ |

Total Score Risk Category

- 0 – 3:** Low risk
- 4 – 7:** Moderate risk
- ≥ 8:** High risk

Webster LR, Webster R. Predicting aberrant behaviors in opioid-treated patients: preliminary validation of the Opioid Risk Tool. Pain Medicine, 2005;6(6):432